

QuikChange Lightning Site-Directed Mutagenesis Kit

Instruction Manual

Catalog # 210518 (10 reactions) and #210519 (30 reactions)

Revision D

Research Use Only. Not for Use in Diagnostic Procedures.

210518-12



LIMITED PRODUCT WARRANTY

This warranty limits our liability to replacement of this product. No other warranties of any kind, express or implied, including without limitation, implied warranties of merchantability or fitness for a particular purpose, are provided by Agilent. Agilent shall have no liability for any direct, indirect, consequential, or incidental damages arising out of the use, the results of use, or the inability to use this product.

ORDERING INFORMATION AND TECHNICAL SERVICES

United States and Canada

Agilent Technologies
Stratagene Products Division

11011 North Torrey Pines Road
La Jolla, CA 92037

Telephone (858) 373-6300
Order Toll Free (800) 424-5444
Technical Services (800) 894-1304
Email techservices@agilent.com
World Wide Web www.genomics.agilent.com

Europe

Location	Telephone
Austria	01 25125 6800
Benelux	02 404 92 22
Denmark	45 70 13 00 30
Finland	010 802 220
France	0810 446 446
Germany	0800 603 1000
Italy	800 012575
Netherlands	020 547 2600
Spain	901 11 68 90
Sweden	08 506 4 8960
Switzerland	0848 8035 60
UK/Ireland	0845 712 5292

All Other Countries

Please contact your local distributor. A complete list of distributors is available at www.genomics.agilent.com.

QuikChange Lightning Site-Directed Mutagenesis Kit

CONTENTS

Materials Provided	1
Storage Conditions	1
Additional Materials Required	1
Introduction	2
QuikChange Lightning Mutagenesis Control.	4
Mutagenic Primer Design	5
Primer Design Guidelines.....	5
Additional Primer Considerations	6
Protocol	7
Mutant Strand Synthesis Reaction (Thermal Cycling).....	7
<i>Dpn</i> I Digestion of the Amplification Products.....	8
Transformation of XL10-Gold Ultracompetent Cells	9
Transformation Guidelines	12
Storage Conditions	12
Aliquoting Cells	12
Use of 14-ml BD Falcon Polypropylene Round-Bottom Tubes.....	12
Use of β -Mercaptoethanol.....	12
Quantity of DNA Added	12
Length and Temperature of the Heat Pulse	12
Preparing the Agar Plates for Color Screening	12
Troubleshooting	13
Preparation of Media and Reagents	15
References	15
MSDS Information	15

QuikChange Lightning Site-Directed Mutagenesis Kit

MATERIALS PROVIDED

Materials Provided	Quantity	
	Catalog #210518 ^a	Catalog #210519 ^b
QuikChange Lightning Enzyme	10 reactions	30 reactions
10× QuikChange Lightning Buffer	500 µl	500 µl
dNTP mix ^{c,d}	10 µl	30 µl
QuikSolution reagent	500 µl	500 µl
Dpn I restriction enzyme ^d	10 reactions	30 reactions
pWhitescript 4.5-kb control plasmid (5 ng/µl)	50 ng	50 ng
Oligonucleotide control primer #1 [34-mer (100 ng/µl)] 5' CCA TGA TTA CGC CAA GCG CGC AAT TAA CCC TCA C 3'	750 ng	750 ng
Oligonucleotide control primer #2 [34-mer (100 ng/µl)] 5' GTG AGG GTT AAT TGC GCG CTT GGC GTA ATC ATG G 3'	750 ng	750 ng
XL10-Gold ultracompetent cells ^e (yellow tubes)	4 × 135 µl	10 × 135 µl
XL10-Gold β-mercaptoethanol mix (β-ME)	50 µl	2 × 50 µl
pUC18 control plasmid (0.1 ng/µl in TE buffer ^f)	10 µl	10 µl

^a The QuikChange Lightning Site-Directed Mutagenesis Kit (Catalog #210518) contains enough reagents for 10 total reactions, which includes 2 control reactions.

^b The QuikChange Lightning Site-Directed Mutagenesis Kit (Catalog #210519) contains enough reagents for 30 total reactions, which includes 2 control reactions.

^c Thaw the dNTP mix once, prepare single-use aliquots, and store the aliquots at -20°C. **Do not subject the dNTP mix to multiple freeze-thaw cycles.**

^d **The composition of the dNTP mix and Dpn I enzyme are proprietary.** These reagents have been optimized for the QuikChange Lightning protocols and have been qualified for use in conjunction with the other kit components. Do not substitute with dNTP mixes or Dpn I enzyme formulations provided with other Stratagene kits or from other sources.

^e Genotype: *Tet*^r*Δ(mcrA)183 Δ(mcrCB-hsdSMR-mrr)173 endA1 supE44 thi-1 recA1 gyrA96 relA1 lac Hte*
[F' proAB *lacI*^q*ZΔM15* *Tn10* (*Tet*^r) *Amp* *Cam*^r]

^f See Preparation of Media and Reagents.

STORAGE CONDITIONS


XL10-Gold Ultracompetent cells, XL10-Gold β-ME, and pUC18 Control Plasmid: -80°C
All Other Components: -20°C

ADDITIONAL MATERIALS REQUIRED

14-ml BD Falcon polypropylene round-bottom tubes (BD Biosciences Catalog #352059)
5-Bromo-4-chloro-3-indolyl-β-D-galactopyranoside (X-gal)
Isopropyl-1-thio-β-D-galactopyranoside (IPTG)

INTRODUCTION

The QuikChange Lightning Site-Directed Mutagenesis Kit* delivers mutant plasmids up to three times faster than our original QuikChange kits, without losses in mutagenesis efficiency or accuracy. The kit has been optimized for mutagenesis of plasmids of up to 14 kb, allowing rapid, efficient, and accurate mutagenesis of small and large plasmids with a single kit. Using the most advanced high fidelity enzyme technology, the protocols have been accelerated while maintaining the highest accuracy for site-directed mutagenesis. Exclusive to the QuikChange Lightning Site-Directed Mutagenesis Kit is a proprietary *Pfu*-based polymerase blend and the newly optimized *Dpn* I enzyme, which together allow for mutagenesis in approximately one hour, plus an overnight transformation.



* Based on a 5-kb plasmid; excludes ramping time.

FIGURE 1 Overview of the QuikChange Lightning site-directed mutagenesis method.

* U.S. Patent Nos. 7,176,004; 7,132,265; 7,045,328; 6,734,293; 6,489,150; 6,444,428; 6,391,548; 6,183,997; 5,948,663; 5,932,419; 5,866,395; 5,789,166; 5,545,552, and patents pending.

In vitro site-directed mutagenesis is an invaluable technique for characterizing the dynamic, complex relationships between protein structure and function, for studying gene expression elements, and for carrying out vector modification. Several approaches to this technique have been published, but these methods generally require single-stranded DNA (ssDNA) as the template^{1–4} and are labor intensive or technically difficult. Our QuikChange Lightning Site-Directed Mutagenesis Kit allows site-specific mutation in virtually any double-stranded plasmid, thus eliminating the need for subcloning and for ssDNA rescue.⁵ In addition, the QuikChange Lightning Site-Directed Mutagenesis Kit does not require specialized vectors, unique restriction sites, multiple transformations or in vitro methylation treatment steps. The simple, rapid three-step procedure requires only about one hour prior to transformation (for plasmids up to 5 kb), and generates mutants with greater than 85% efficiency in a single reaction (see Figure 1).

The QuikChange Lightning Enzyme is a novel proprietary blend that includes a derivative of *PfuUltra* high-fidelity (HF) DNA polymerase** for mutagenic primer-directed replication of both plasmid strands with the highest fidelity.⁶ The basic procedure utilizes a supercoiled double-stranded DNA (dsDNA) vector with an insert of interest and two synthetic oligonucleotide primers, both containing the desired mutation (see Figure 1). The oligonucleotide primers, each complementary to opposite strands of the vector, are extended during temperature cycling by *PfuUltra* HF DNA polymerase, without primer displacement. Extension of the oligonucleotide primers generates a mutated plasmid containing staggered nicks. Following temperature cycling, the product is treated with *Dpn* I. The *Dpn* I endonuclease (target sequence: 5'-Gm⁶ATC-3') is specific for methylated and hemimethylated DNA and is used to digest the parental DNA template and to select for mutation-containing synthesized DNA.⁶ (DNA isolated from almost all *E. coli* strains is dam methylated and therefore susceptible to *Dpn* I digestion.) The nicked vector DNA containing the desired mutations is then transformed into XL10-Gold ultracompetent cells.

Note While plasmid DNA isolated from almost all of the commonly used *E. coli* strains (dam⁺) is methylated and is a suitable template for mutagenesis, plasmid DNA isolated from the exceptional dam⁻ *E. coli* strains, including JM110 and SCS110, is not suitable.

Unwanted second-site errors are virtually eliminated and high mutation efficiencies are obtained using this method due to the high fidelity of the *PfuUltra* HF DNA polymerase, the use of a small amount of starting DNA template and the use of a low number of thermal cycles.

The QuikChange Lightning Site-Directed Mutagenesis Kit may be used to make point mutations, replace amino acids, and delete or insert single or multiple adjacent amino acids. The kit has been optimized with plasmids ranging in size from 4–14 kb.

** U.S. Patent Nos. 7,045,328; 6,734,293; 6,489,150; 6,444,428; 6,183,997; 5,948,663; 5,866,395; 5,545,552 and patents pending.

⁶ *PfuUltra* HF DNA polymerase has 18-fold higher fidelity in DNA synthesis than *Taq* DNA polymerase.

QUIKCHANGE LIGHTNING MUTAGENESIS CONTROL

The pWhitescript 4.5-kb control plasmid is used to test the efficiency of mutant plasmid generation using the QuikChange Lightning Site-Directed Mutagenesis Kit. The pWhitescript 4.5-kb control plasmid contains a stop codon (TAA) at the position where a glutamine codon (CAA) would normally appear in the β -galactosidase gene of the pBluescript II SK(–) phagemid (corresponding to amino acid 9 of the protein). XL10-Gold ultracompetent cells transformed with this control plasmid appear white on LB–ampicillin agar plates (see *Preparation of Media and Reagents*), containing IPTG and X-gal, because β -galactosidase activity has been obliterated. The oligonucleotide control primers create a point mutation on the pWhitescript 4.5-kb control plasmid that reverts the T residue of the stop codon (TAA) at amino acid 9 of the β -galactosidase gene to a C residue, to produce the glutamine codon (CAA) found in the wild-type sequence. Following transformation, colonies can be screened for the β -galactosidase (β -gal $^{+}$) phenotype of blue color on media containing IPTG and X-gal.

MUTAGENIC PRIMER DESIGN

Note *Mutagenic primers can be designed using our web-based QuikChange Primer Design Program available online at www.agilent.com/genomics/qcpd.*

Primer Design Guidelines

The mutagenic oligonucleotide primers for use in this protocol must be designed individually according to the desired mutation. The following considerations should be made when designing mutagenic primers:

- ◆ Both of the mutagenic primers must contain the desired mutation and anneal to the same sequence on opposite strands of the plasmid.
- ◆ Primers should be between 25 and 45 bases in length, with a melting temperature (T_m) of $\geq 78^\circ\text{C}$. Primers longer than 45 bases may be used, but using longer primers increases the likelihood of secondary structure formation, which may affect the efficiency of the mutagenesis reaction.
- ◆ The following formula is commonly used for estimating the T_m of primers:

$$T_m = 81.5 + 0.41(\%GC) - (675/N) - \% \text{ mismatch}$$

For calculating T_m :

- N is the primer length in bases
- values for **%GC** and **% mismatch** are whole numbers

For calculating T_m for primers intended to introduce insertions or deletions, use this modified version of the above formula:

$$T_m = 81.5 + 0.41(\%GC) - (675/N)$$

where N does not include the bases which are being inserted or deleted.

Note *When using primer design software for QuikChange Lightning site-directed mutagenesis applications, be aware that the T_m calculated by the primer design software may differ from the T_m value calculated using the formula presented above. we recommend verifying primer T_m 's using the formula above or by using the QuikChange T_m calculator, available online at www.genomics.agilent.com.*

- ◆ The desired mutation (deletion or insertion) should be in the middle of the primer with $\sim 10\text{--}15$ bases of correct sequence on both sides.
- ◆ The primers optimally should have a minimum GC content of 40% and should terminate in one or more C or G bases.
- ◆ Primers need not be 5' phosphorylated.

- ◆ For typical mutagenic primers, purification by desalting is generally sufficient. For long or complex mutagenic primers, purification either by liquid chromatography (HPLC/FPLC) or by polyacrylamide gel electrophoresis (PAGE) may produce a significant increase in mutagenesis efficiency.

Additional Primer Considerations

- ◆ The mutagenesis protocol uses 125 ng of each oligonucleotide primer. To convert nanograms to picomoles of oligo, use the following equation:

$$X \text{ pmoles of oligo} = \frac{\text{ng of oligo}}{330 \times \# \text{ of bases in oligo}} \times 1000$$

For example, for 125 ng of a 25-mer:

$$\frac{125 \text{ ng of oligo}}{330 \times 25 \text{ bases}} \times 1000 = 15 \text{ pmole}$$

- ◆ It is important to keep primer concentration in excess. We suggest varying the amount of template while keeping the concentration of the primer constantly in excess.

PROTOCOL

Mutant Strand Synthesis Reaction (Thermal Cycling)

Notes Ensure that the plasmid DNA template is isolated from a dam⁺ E. coli strain. The majority of the commonly used E. coli strains are dam⁺. Plasmid DNA isolated from dam⁻ strains (e.g. JM110 and SCS110) is not suitable.

To maximize temperature-cycling performance, we strongly recommend using thin-walled tubes, which ensure ideal contact with the temperature cycler's heat blocks. The following protocols were optimized using thin-walled tubes.

1. Synthesize two complimentary oligonucleotides containing the desired mutation, flanked by unmodified nucleotide sequence. Purify these oligonucleotide primers prior to use in the following steps (see *Mutagenic Primer Design*).

2. Prepare the control reaction as indicated below:

5 µl of 10× reaction buffer
5 µl (25 ng) of pWhitescript 4.5-kb control plasmid (5 ng/µl)
1.25 µl (125 ng) of oligonucleotide control primer #1
[34-mer (100 ng/µl)]
1.25 µl (125 ng) of oligonucleotide control primer #2
[34-mer (100 ng/µl)]
1 µl of dNTP mix
1.5 µl of QuikSolution reagent
34 µl ddH₂O (to bring the final reaction volume to 50 µl)

Then add:

1 µl of QuikChange Lightning Enzyme

3. Prepare the sample reaction(s) as indicated below:

Note Set up a series of sample reactions using various amounts of dsDNA template ranging from 10 to 100 ng (e.g., 10, 25, 50, and 100 ng of dsDNA template) while keeping the primer concentration constant.

5 µl of 10× reaction buffer
 X µl (10–100 ng) of dsDNA template
 X µl (125 ng) of oligonucleotide primer #1
 X µl (125 ng) of oligonucleotide primer #2
1 µl of dNTP mix
1.5 µl of QuikSolution reagent
ddH₂O to a final volume of 50 µl

Then add:

1 µl of QuikChange Lightning Enzyme

4. If the thermal cycler to be used does not have a hot-top assembly, overlay each reaction with ~30 µl of mineral oil.
5. Cycle each reaction using the cycling parameters outlined in Table I. (For the control reaction, use a 2.5-minute extension time.)

TABLE I

Cycling Parameters for the QuikChange Lightning Site-Directed Mutagenesis Method

Segment	Cycles	Temperature	Time
1	1	95°C	2 minutes
2	18	95°C	20 seconds
		60°C	10 seconds
		68°C	30 seconds/kb of plasmid length*
3	1	68°C	5 minutes

* For example, a 5-kb plasmid requires 2.5 minutes per cycle at 68°C.

Dpn I Digestion of the Amplification Products

1. Add 2 µl of the provided *Dpn* I restriction enzyme directly to each amplification reaction.

Notes Use only the *Dpn* I enzyme provided; do not substitute with an enzyme from another source.

*If a mineral oil overlay was used during the synthesis reaction, it is important to insert the pipet tip below the overlay when adding the *Dpn* I enzyme to the reaction tubes and when transferring the 2 µl of *Dpn* I-treated DNA to the transformation reaction. Using specialized aerosol-resistant pipet tips, which are small and pointed, facilitates this process.*

2. Gently and thoroughly mix each reaction mixture by pipetting the solution up and down several times. Briefly spin down the reaction mixtures and then immediately incubate at 37°C for 5 minutes to digest the parental (i.e., the nonmutated) supercoiled dsDNA.

Transformation of XL10-Gold Ultracompetent Cells

Notes Please read the Transformation Guidelines before proceeding with the transformation protocol.

XL10-Gold cells are resistant to tetracycline and chloramphenicol. If the mutagenized plasmid contains only the *tet^R* or *cam^R* resistance marker, an alternative strain of competent cells must be used.

1. Gently thaw the XL10-Gold ultracompetent cells on ice. For each control and sample reaction to be transformed, aliquot 45 µl of the ultracompetent cells to a *prechilled* 14-ml BD Falcon polypropylene round-bottom tube.
2. Add 2 µl of the β-ME mix provided with the kit to the 45 µl of cells. (Using an alternative source of β-ME may reduce transformation efficiency.)
3. Swirl the contents of the tube gently. Incubate the cells on ice for 2 minutes.
4. Transfer 2 µl of the *Dpn* I-treated DNA from each control and sample reaction to separate aliquots of the ultracompetent cells.

Note If mineral oil overlays were used during thermal cycling, carefully remove any residual mineral oil from the pipet tip before transferring the *Dpn* I-treated DNA to the transformation reaction.

As an optional control, verify the transformation efficiency of the XL10-Gold ultracompetent cells by adding 1 µl of 0.01 ng/µl pUC18 control plasmid (dilute the control provided 1:10 in high-quality water) to another 45-µl aliquot of cells.

5. Swirl the transformation reactions gently to mix and incubate the reactions on ice for 30 minutes.

Note The incubation time for this step may be reduced to 10 minutes without substantial losses in transformation efficiency.

6. Preheat NZY⁺ broth (see Preparation of Media and Reagents) in a 42°C water bath for use in step 9.

Note Transformation of XL10-Gold ultracompetent cells has been optimized using NZY⁺ broth.

7. Heat-pulse the tubes in a 42°C water bath for 30 seconds. The duration of the heat pulse is *critical* for obtaining the highest efficiencies. Do not exceed 42°C.

Note *This heat pulse has been optimized for transformation in 14-ml BD Falcon polypropylene round-bottom tubes.*

8. Incubate the tubes on ice for 2 minutes.
9. Add 0.5 ml of preheated (42°C) NZY⁺ broth to each tube, then incubate the tubes at 37°C for 1 hour with shaking at 225–250 rpm.
10. Plate the appropriate volume of each transformation reaction, as indicated in the table below, on agar plates containing the appropriate antibiotic for the plasmid vector.

For the mutagenesis and transformation controls, spread cells on LB–ampicillin agar plates containing 80 µg/ml X-gal and 20 mM IPTG (see *Preparing the Agar Plates for Color Screening*).

Transformation reaction plating volumes

Reaction Type	Volume to Plate ^a
pWhitescrit mutagenesis control	10 µl
pUC18 transformation control	2.5 µl
Sample mutagenesis	10–250 µl ^b

^a When plating volumes less than 100 µl, place a 200-µl pool of NZY⁺ broth on the agar plate, pipet the small volume of the transformation reaction into the pool, then spread the mixture.

^b The optimal amount for spreading varies according to the size and sequence of the mutagenized plasmid. It is generally useful to plate the entire transformation mixture, divided among multiple plates and covering a range of plating volumes.

11. Incubate the transformation plates at 37°C for >16 hours.

Expected Results for the Control Transformations

The expected colony number from the transformation of the pWhitescrit 4.5 kb control mutagenesis reaction is >100 colonies. Greater than 85% of the colonies should contain the mutation and appear as blue colonies on agar plates containing IPTG and X-gal.

Note *The mutagenesis efficiency (ME) for the pWhitescrit 4.5-kb control plasmid is calculated by the following formula:*

$$ME = \frac{\text{Number of blue colony forming units (cfu)}}{\text{Total number of colony forming units (cfu)}} \times 100\%$$

If transformation of the pUC18 control plasmid was performed, >50 colonies (>10⁹ cfu/µg) should be observed, with >98% having the blue phenotype.

Expected Results for Sample Transformations

The expected colony number depends upon the base composition and length of the DNA template employed. For suggestions on increasing colony number, see *Troubleshooting*. The insert of interest should be sequenced to verify that selected clones contain the desired mutation(s).

TRANSFORMATION GUIDELINES

Storage Conditions

Ultracompetent cells are sensitive to even small variations in temperature and must be stored at the bottom of a -80°C freezer. Transferring tubes from one freezer to another may result in a loss of efficiency. Ultracompetent cells should be placed at -80°C directly from the dry ice shipping container.

Aliquoting Cells

When aliquoting, keep ultracompetent cells on ice at all times. It is essential that the BD Falcon polypropylene tubes are placed on ice before the cells are thawed and that the cells are aliquoted directly into the prechilled tubes.

Use of 14-ml BD Falcon Polypropylene Round-Bottom Tubes

It is important that 14-ml BD Falcon polypropylene round-bottom tubes (BD Biosciences Catalog #352059) are used for the transformation protocol, since other tubes may be degraded by the β -mercaptoethanol used in the *Transformation Protocol*. In addition, the duration of the heat-pulse step is critical and has been optimized specifically for the thickness and shape of these tubes.

Use of β -Mercaptoethanol

β -Mercaptoethanol (β -ME) has been shown to increase transformation efficiency. The XL10-Gold β -mercaptoethanol mix provided in this kit is diluted and ready to use.

Quantity of DNA Added

Greatest efficiencies are observed when adding 2 μ l of the synthesis reaction. A greater number of colonies will be obtained when adding a greater volume of the synthesis reaction, although the overall efficiency may be lower.

Length and Temperature of the Heat Pulse

There is a defined window of highest efficiency resulting from the heat pulse during transformation. Optimal efficiencies are observed when cells are heat-pulsed for 30 seconds. Do not exceed 42°C.

Preparing the Agar Plates for Color Screening

To prepare the LB agar plates for blue-white color screening, add 80 μ g/ml of 5-bromo-4-chloro-3-indolyl- β -D-galactopyranoside (X-gal), 20 mM isopropyl-1-thio- β -D-galactopyranoside (IPTG), and the appropriate antibiotic to the LB agar. Alternatively, 100 μ l of 10 mM IPTG and 100 μ l of 2% X-gal can be spread on the LB agar plates 30 minutes prior to plating the transformations. Prepare the IPTG in sterile dH₂O; prepare the X-gal in dimethylformamide (DMF). Do not mix the IPTG and X-gal before pipetting them onto the plates because these chemicals may precipitate.

TROUBLESHOOTING

When used according to the guidelines outlined in this instruction manual, this kit provides a reliable means to conduct site-directed mutagenesis using dsDNA templates. Variations in the base composition and length of the DNA template and in thermal cycler performance may contribute to differences in mutagenesis efficiency. We provide the following guidelines for troubleshooting these variations.

Observation	Suggestion(s)
Low transformation efficiency or low colony number	If using a mineral oil overlay during thermal cycling, ensure that excess mineral oil is not transferred into the transformation reaction when pipetting the <i>Dpn</i> I-treated DNA. Using the smallest pipet tips available, insert the pipet tip completely below the mineral layer overlay and clear the pipet tip while submerged beneath the mineral oil overlay before collecting the sample.
	Ensure that sufficient mutant DNA is synthesized in the reaction. Increase the amount of the <i>Dpn</i> I-treated DNA used in the transformation reaction to 3 µl. If a greater number of colonies is desired, the transformation reaction may be directly scaled up to include 90 µl of competent cells and 4–6 µl of the <i>Dpn</i> I-treated DNA.
	Visualize the DNA template on a gel to verify the quantity and quality. Nicked or linearized plasmid DNA will not generate complete circular product. Verify that the template DNA is at least 80% supercoiled.
	It is not uncommon to observe low numbers of colonies, especially when generating large mutations. Most of the colonies that do appear, however, will contain mutagenized plasmid.
	Ethanol precipitate the <i>Dpn</i> I-digested PCR product, and resuspend in a decreased volume of water before transformation.
Low mutagenesis efficiency or low colony number with the control reaction	Ensure that the 10× QuikChange Lightning Buffer provided with this kit was used for the synthesis reaction. The QuikChange Lightning protocols have been optimized specifically with this buffer; substitution with another buffer system results in decreased mutagenesis efficiency and/or fidelity.
	Ensure that the <i>Dpn</i> I enzyme provided with this kit was used for the digestion reaction. Substitution with another enzyme formulation will decrease mutagenesis efficiency.
	Different thermal cyclers may contribute to variations in amplification efficiencies. Optimize the cycling parameters for the control reaction and then repeat the protocol for the sample reactions using the adjusted parameters.
	Ensure that supercompetent cells are stored at the bottom of a –80°C freezer immediately upon arrival (see also <i>Transformation Guidelines</i>).
	Verify that the agar plates were prepared correctly. See <i>Preparing the Agar Plates for Color Screening</i> , and follow the recommendations for IPTG and X-Gal concentrations carefully.
	For best visualization of the blue (β -gal ⁺) phenotype, the control plates must be incubated for at least 16 hours at 37°C.
	Avoid multiple freeze-thaw cycles for the dNTP mix. Thaw the dNTP mix once, prepare single-use aliquots, and store the aliquots at –20°C. Do not subject the dNTP mix to multiple freeze-thaw cycles.

Table continues on following page

Table continues from the previous page

Low mutagenesis efficiency with the sample reaction(s)	<p>Ensure that the <i>Dpn</i> I enzyme provided with this kit was used for the digestion reaction. Substitution with another enzyme formulation will decrease mutagenesis efficiency.</p> <p>If using a mineral oil overlay add the <i>Dpn</i> I restriction enzyme below the mineral oil in the digestion step and ensure proper mixing of all reaction components, especially the <i>Dpn</i> I.</p> <p>If excess DNA template was present, it may be beneficial to increase the incubation time for the <i>Dpn</i> I digestion, to ensure complete digestion of the parental template.</p> <p>Avoid multiple freeze-thaw cycles for the dNTP mix. Thaw the dNTP mix once, prepare single-use aliquots, and store the aliquots at –20°C. Do not subject the dNTP mix to multiple freeze-thaw cycles.</p> <p>The formation of secondary structures may be inhibiting the mutagenesis reaction. Increasing the annealing temperature up to 68°C may help to alleviate secondary structure formation and improve mutagenesis efficiency.</p>
False positives	<p>Poor quality primers can lead to false positives. Radiolabel the primers and check for degradation on an acrylamide gel. Resynthesize the primers. If primers were used without purification, purify primers by liquid chromatography or PAGE.</p> <p>False priming can lead to false positives. Increase the stringency of the reaction by increasing the annealing temperature up to 68°C.</p>
Absence of amplification product when analyzed by gel electrophoresis	<p>Although it is not part of the standard protocol, some researchers choose to verify amplification by gel electrophoresis prior to transformation (typically, 10 µl of the synthesis product is analyzed on a 1% agarose gel). While a positive result verifies successful synthesis, a negative result does not indicate failure of the synthesis reaction. Reactions that produce sufficient numbers of mutant plasmid colonies after transformation may or may not produce a visible band at this stage. We recommend proceeding with <i>Dpn</i> I digestion and transformation when gel electrophoresis analysis yields negative results.</p>

PREPARATION OF MEDIA AND REAGENTS

LB Agar (per Liter) 10 g of NaCl 10 g of tryptone 5 g of yeast extract 20 g of agar Add deionized H ₂ O to a final volume of 1 liter Adjust pH to 7.0 with 5 N NaOH Autoclave Pour into petri dishes (~25 ml/100-mm plate)	LB-Ampicillin Agar (per Liter) (Use for reduced satellite colony formation) 1 liter of LB agar Autoclave Cool to 55°C Add 100 mg of filter-sterilized ampicillin Pour into petri dishes (~25 ml/100-mm plate)
NZY⁺ Broth (per Liter) 10 g of NZ amine (casein hydrolysate) 5 g of yeast extract 5 g of NaCl Add deionized H ₂ O to a final volume of 1 liter Adjust to pH 7.5 using NaOH Autoclave Add the following filter-sterilized supplements prior to use: 12.5 ml of 1 M MgCl ₂ 12.5 ml of 1 M MgSO ₄ 20 ml of 20% (w/v) glucose (or 10 ml of 2 M glucose)	TE Buffer 10 mM Tris-HCl (pH 7.5) 1 mM EDTA

REFERENCES

1. Kunkel, T. A. (1985) *Proc Natl Acad Sci U S A* 82(2):488-92.
2. Sugimoto, M., Esaki, N., Tanaka, H. and Soda, K. (1989) *Anal Biochem* 179(2):309-11.
3. Taylor, J. W., Ott, J. and Eckstein, F. (1985) *Nucleic Acids Res* 13(24):8765-85.
4. Vandeyar, M. A., Weiner, M. P., Hutton, C. J. and Batt, C. A. (1988) *Gene* 65(1):129-33.
5. Papworth, C., Bauer, J. C., Braman, J. and Wright, D. A. (1996) *Strategies* 9(3):3-4.
6. Nelson, M. and McClelland, M. (1992) *Methods Enzymol* 216:279-303.

MSDS INFORMATION

The Material Safety Data Sheet (MSDS) information for Stratagene products is provided on the web at <http://www.genomics.agilent.com>. MSDS documents are not included with product shipments.

STRATAGENE

An Agilent Technologies Division

QuikChange Lightning Site-Directed Mutagenesis Kit

Catalog #210518 and #210519

QUICK-REFERENCE PROTOCOL

- ◆ Prepare the control and sample reaction(s) as indicated below:

Note Set up a series of sample reactions using various amounts of dsDNA template (e.g., 10, 25, 50, and 100 ng of dsDNA template).

Control Reaction

5 µl of 10× reaction buffer
5 µl (25 ng) of pWhitescript 4.5-kb control template (5 ng/µl)
1.25 µl (125 ng) of control primer #1
1.25 µl (125 ng) of control primer #2
1 µl of dNTP mix
1.5 µl of QuikSolution reagent
34 µl ddH₂O (for a final volume of 50 µl)

Sample Reaction

5 µl of 10× reaction buffer
X µl (10–100 ng) of dsDNA template
X µl (125 ng) of oligonucleotide primer #1
X µl (125 ng) of oligonucleotide primer #2
1 µl of dNTP mix
1.5 µl of QuikSolution reagent
ddH₂O to a final volume of 50 µl

- ◆ Add 1 µl of QuikChange Lightning Enzyme to each control and sample reaction

Note If the thermal cycler does not have a hot-top, overlay each reaction with ~30 µl of mineral oil.

- ◆ Cycle each reaction using the cycling parameters outlined in the following table:

Segment	Cycles	Temperature	Time
1	1	95°C	2 minutes
2	18	95°C	20 seconds
		60°C	10 seconds
		68°C	30 seconds /kb of plasmid length
3	1	68°C	5 minutes

- ◆ Add 2 µl of the Dpn I restriction enzyme (below the mineral oil overlay, if used)
- ◆ Gently and thoroughly mix each reaction, microcentrifuge briefly, then immediately incubate at 37°C for 5 minutes to digest the parental dsDNA
- ◆ Transform 2 µl of the Dpn I-treated DNA from each reaction into separate 45-µl aliquots of XL10-Gold ultracompetent cells (see Transformation of XL10-Gold Ultracompetent Cells in the instruction manual)